



<https://printo.it/pediatric-rheumatology/TW/intro>

## MDK (IgD)

### 1. MDK

#### 1.1 MDK

MDK (IgD) is a marker of disease activity in children with systemic lupus erythematosus (SLE). It is a type of immunoglobulin (Ig) that is produced by B cells. MDK is a dimeric protein, meaning it consists of two identical subunits. The presence of MDK in the blood is associated with active disease, particularly in the form of skin rashes and joint pain. MDK levels are measured in the blood and are used to monitor disease activity and response to treatment.

#### 1.2 MDK

MDK levels are measured in the blood and are used to monitor disease activity and response to treatment. MDK levels are measured in the blood and are used to monitor disease activity and response to treatment. MDK levels are measured in the blood and are used to monitor disease activity and response to treatment.

#### 1.3 MDK

MDK is a marker of disease activity in children with systemic lupus erythematosus (SLE). It is a type of immunoglobulin (Ig) that is produced by B cells. MDK is a dimeric protein, meaning it consists of two identical subunits. The presence of MDK in the blood is associated with active disease, particularly in the form of skin rashes and joint pain. MDK levels are measured in the blood and are used to monitor disease activity and response to treatment.

#### 1.4 MDK

MDK levels are measured in the blood and are used to monitor disease activity and response to treatment. MDK levels are measured in the blood and are used to monitor disease activity and response to treatment. MDK levels are measured in the blood and are used to monitor disease activity and response to treatment.

#### 1.5 MDK

MDK levels are measured in the blood and are used to monitor disease activity and response to treatment. MDK levels are measured in the blood and are used to monitor disease activity and response to treatment. MDK levels are measured in the blood and are used to monitor disease activity and response to treatment.

